

(revision date:2/14/2019)

Pepper, Eggplant: Flea beetles

Use *Integrated Pest Management (IPM)* for successful plant problem management.

Biology

Flea beetles are small, brown to metallic black beetles with a habit of jumping like fleas when disturbed. They feed on many plants including beets, kale, collards, radish, and many weeds, particularly those in the mustard family. Small round holes are eaten in the leaves, usually early in the season. The larvae typically feed on underground portions of the host plants. Damage can be very severe on seedling plants.

Management Options

Non-Chemical Management

~ Keep garden and adjacent areas clear of weeds which may attract flea beetles.

Select non-chemical management options as your first choice!

Chemical Management

IMPORTANT: Visit *Home and Garden Fact Sheets* for more information on using pesticides


Apply according to label instructions when flea beetles are first seen.

Listed below are examples of pesticides that are legal in Washington. Always read and follow all label directions.

- ~ Bonide Captain Jack's Deadbug Brew R-T-U [Organic]
 - active ingredient: *spinosad (spinosyn A+D)*
 - EPA reg no: 4-472
 - ~ Bonide Eight Insect Control Garden & Home R-T-U
 - active ingredient: *permethrin*
 - EPA reg no: 4-406
 - ~ Bug Buster-O [Organic]
 - active ingredient: *pyrethrins*
 - EPA reg no: 1021-1771-54705
 - ~ Monterey Garden Insect Spray [Organic]
 - active ingredient: *spinosad (spinosyn A+D)*
 - EPA reg no: 62719-314-54705
 - ~ Naturalis L
 - active ingredient: *Beauveria bassiana*
 - EPA reg no: 53871-9
 - ~ Safer Brand BioNEEM Multi-Purpose Insecticide & Repellent Conc [Organic]
 - active ingredient: *azadirachtin*
 - EPA reg no: 70051-6-42697
- ~ This list may not include all products registered for this use.

Pepper, Eggplant: Flea beetles

Images



~ Caption: Adult flea beetle damage on potato leaf
~ Photo by: Unknown



~ Caption: Adult flea beetle damage on potatoes
~ Photo by: S.J. Collman