

(revision date:5/1/2013)

Herbicide Damage: Dichlobenil (Casoron)

Use Integrated Pest Management (IPM) for successful plant problem management.

Biology

Dichlobenil is a soil-applied herbicide which inhibits root development and prevents germination and establishment of weed seedlings. While dichlobenil is not translocated in the plant, symptoms resulting from inhibited root growth may appear on aboveground portions of the plant. The primary aboveground symptom of damage is chlorosis. On broad-leaved plants this can be tip, marginal, or interveinal yellowing, or can appear as overall yellowing of the leaf. On conifers, excessive dichlobenil applications can cause tip chlorosis tending toward tip necrosis. Symptoms usually appear on the new growth. The damage may occur on those leaves and leaf parts that orient toward the afternoon sun. Dichlobenil damage closely resembles triazine/simazine damage. This herbicide is persistent in the soil and decomposes slowly.

Management Options

Non-Chemical Management

- ~ Do not overapply or apply dichlobenil near sensitive plants.
- ~ Read pesticide labels carefully prior to purchase and application.

Select non-chemical management options as your first choice!

Chemical Management

IMPORTANT: Visit Home and Garden Fact Sheets for more information on using pesticides

Carefully read all label instructions prior to using products containing dichlobenil.

Herbicide Damage: Dichlobenil (Casoron)

Images




~ Caption: Pine casoron damage
~ Photo by: R. Maleike



~ Caption: Rhododendron casoron damage
~ Photo by: R. Maleike



~ Caption: Turf casoron damage
~ Photo by: R. Maleike



~ Caption: European hornbeam casoron damage
~ Photo by: R. Maleike



~ Caption: Land firethorn casoron damage
~ Photo by: R. Maleike



~ Caption: Kwanzan cherry casoron damage
~ Photo by: R. Maleike